

## Would be IL-6 a missing link between chronic inflammatory rheumatism and depression after chikungunya infection?

Alfonso J. Rodriguez-Morales<sup>1</sup>  · Karol Liceth Hoyos-Guapacha<sup>1</sup> ·  
Sara Lucia Vargas-Zapata<sup>1</sup> · Oscar Mauricio Meneses-Quintero<sup>1</sup> ·  
Julio César Gutiérrez-Segura<sup>1</sup>

Received: 11 April 2017 / Accepted: 17 May 2017 / Published online: 26 May 2017  
© Springer-Verlag Berlin Heidelberg 2017

Dear Sir,

As has been previously reported [1], chikungunya virus disease (CHIK) has emerged in Latin America as a significant acute infectious disease condition, but also with multiple implications during its chronic phase, including the post-chikungunya chronic inflammatory rheumatism (pCHIK-CIR), now well-documented, particularly in Colombia, where around 50% of patients go with clinical rheumatological, but even systemic disease, beyond 12 weeks after infection (chronic phase) [2, 3]. Among the systemic reported conditions, depression has been described in some studies [4].

Major depressive disorder is a severe mental illness. It causes disability and even mortality worldwide [5, 6]. Depression is a remarkable problem, with relevant direct and indirect costs, being considered a major public health issue [5]. The pathogenesis of this disorder is still unknown, but there are some evidences that range from several factors or theories: cognitive, hypothalamic–pituitary–adrenal axis dysfunction, neurodegenerative disorders, stress generation, social, psychological and environmental issues, as well inflammation, among others [5].

Among the last of them, depressive symptoms have been linked to an immune response [5–7]. This consideration is based on the highly interrelated extensive relationships between the immune, endocrine and central nervous (CNS) systems.

During periods of increased stress or chronic illness, pro-inflammatory cytokines are released. Chronic exposure to cortisol and cytokines reduces the availability of monoamines, such as serotonin, dopamine and norepinephrine, by influencing synthesis and reuptake. These monoamines play an important role in emotion regulation [5].

The hippocampal and prefrontal cortex are widely associated with the occurrence of mood disorders. The hippocampus is sensitive to cortisol toxicity and elevated pro-inflammatory cytokines; reduction of hippocampal volume has a negative effect on affective modulation also [5]. Neuroinflammation activation leads also to abnormal function in the prefrontal cortex [6].


Among the involved cytokines, interleukin-6 (IL-6) plays an important role in the immune response and acute phase reactions. It is a multifunctional cytokine that regulates the growth and differentiation of various tissues [6]. This cytokine is a versatile protein primarily categorized as proinflammatory but also anti-inflammatory one. It can be released from several tissues including white blood, endothelial and epithelial cells, astrocytes, adipose tissue, microglia and neurons. IL-6 contributes to stress sensitivity in the CNS. IL-6 is considered a relevant biomarker for depression, and it has a potential therapeutic target for mood-related disorders [7].

During acute and chronic chikungunya infection, cytokines, such as IL-6, have been shown to play important roles, as has been also reported in other arthritides, including epidemic polyarthritis caused by Ross River virus and rheumatoid arthritis [8]. At the chronic phase, levels of IL-6 were found to be associated with persistent arthralgia providing a possible explanation for the etiology of arthralgia that affects numerous CHIKV-infected patients [9]. Even more, recently, high scores at the disease activity index 28 (DAS-28) and World

✉ Alfonso J. Rodriguez-Morales  
arodriguezm@utp.edu.co

<sup>1</sup> Research Group Public Health and Infection, Faculty of Health Sciences, Universidad Tecnológica de Pereira, Pereira 660001, Risaralda, Colombia

**Fig. 1** Proposed model of relationships between acute and chronic infection and depression mediated by IL-6 in chikungunya



Health Organization Disablement Assessment Schedule II (WHODAS-II) questionnaires (used to evaluate the clinical evolution of patients with CHIKV infection) and high serum levels of IL-6 in those with acute infection may be predictors for developing subacute and chronic disease [10]. Then, seems clear that the question, would be if IL-6 is a missing link between chronic inflammatory rheumatism and depression after chikungunya infection? Solving the puzzle, during acute CHIKV infection, levels of IL-6 and other cytokines increase (Fig. 1); this would be one of the biological factors related to stress sensitivity together with other social factors that would lead to post-CHIKV depression. Also, after the 12-week period, moving to the chronic phase, in those presenting clinical disease of pCHIK-CIR, IL-6 is also increased (Fig. 1), and this would be also during chronic phase a risk for depression after the acute stage of CHIKV infection.

Given these considerations, IL-6 would be used in the clinical setting to predict pCHIK-CIR, as suggested in previous reports [10], but its assessment in those patients with post-CHIKV depression is also necessary. This would provide plausible mechanisms in the complex area of cytokine signaling that could explain the pathogenesis of CHIK chronic consequences [9], as well in the development of interventions targeting the IL-6 signaling blockade [8] in order to potentially impact on depression and pCHIK-CIR.

In recent years, the relationship between inflammatory biomarkers and CNS disorders, including depression, has been extensively studied; nevertheless, our hypothesis deserves further study [11]. Also, other inflammatory cytokines, including TNF- $\alpha$ , IFN- $\gamma$ , IL-8 and IL-1 $\beta$ , coexist in a variety of clinical disorders. But, associations, such as proposed by us, remain to be elucidated. Further studies are needed to establish causality between inflammation in pCHIK-CIR and depressive symptoms.

#### Compliance with ethical standards

**Conflict of interest** No conflict of interest to declare.

#### References

1. Alfaro-Tolosa P, Clouet-Huerta DE, Rodriguez-Morales AJ (2015) Chikungunya, the emerging migratory rheumatism. *Lancet Infect Dis* 15(5):510–512
2. Rodriguez-Morales AJ, Cardona-Ospina JA, Fernanda Urbano-Garzon S, Sebastian Hurtado-Zapata J (2016) Prevalence of post-chikungunya infection chronic inflammatory arthritis: a systematic review and meta-analysis. *Arthritis Care Res* 68(12):1849–1858
3. Rodriguez-Morales AJ, Gil-Restrepo AF, Ramirez-Jaramillo V et al (2016) Post-chikungunya chronic inflammatory rheumatism: results from a retrospective follow-up study of 283 adult and child cases in La Virginia, Risaralda, Colombia. *F1000Research* 5:360
4. Bhatia MS, Gautam P, Jhanjee A (2015) Psychiatric morbidity in patients with chikungunya fever: first report from India. *J Clin Diagn Res* 9(10):V01–V03
5. Wittenborn AK, Rahmandad H, Rick J, Hosseinichimeh N (2016) Depression as a systemic syndrome: mapping the feedback loops of major depressive disorder. *Psychol Med* 46(3):551–562
6. Zhang C, Wu Z, Zhao G, Wang F, Fang Y (2016) Identification of IL6 as a susceptibility gene for major depressive disorder. *Sci Rep* 6:31264
7. Hodes GE, Menard C, Russo SJ (2016) Integrating Interleukin-6 into depression diagnosis and treatment. *Neurobiol Stress* 4:15–22
8. Chaitanya IK, Muruganandam N, Sundaram SG et al (2011) Role of proinflammatory cytokines and chemokines in chronic arthropathy in CHIKV infection. *Viral Immunol* 24(4):265–271
9. Chow A, Her Z, Ong EK et al (2011) Persistent arthralgia induced by Chikungunya virus infection is associated with interleukin-6 and granulocyte macrophage colony-stimulating factor. *J Infect Dis* 203(2):149–157

10. Sepulveda-Delgado J, Vera-Lastra OL, Trujillo-Murillo K et al (2017) Inflammatory biomarkers, disease activity index, and self-reported disability may be predictors of chronic arthritis after chikungunya infection: brief report. *Clin Rheumatol* 36(3):695–699
11. Kim JW, Szigethy EM, Melhem NM, Saghaifi EM, Brent DA (2014) Inflammatory markers and the pathogenesis of pediatric depression and suicide: a systematic review of the literature. *J Clin Psychiatry* 75(11):1242–1253